

AUFMERKSAM UND GEDULDIG – TEIL 2: BEGLEITUNG DURCH DIE HEBAMME

Hebammenforum 4/2021; 22: 42–46

- 25 Dahlen HG, Homer CS, Leap N, Tracy SK: From social to surgical: historical perspectives on perineal care during labour and birth. *Women Birth* 2011; 24: 105–111
- 26 Altaweli R, McCourt C, Scamell M, Curtis Tyler K: Ethnographic study of the use of interventions during the second stage of labor in Jeddah, Saudi Arabia. *Birth* 2019; 46: 500–508
- 27 Jesacher K: Kritik an strikter Anleitung zu forciertem Pressen und Rückenlage der Gebärenden. *Hebamme* 2016; 29: 160–165
- 28 Hollins Martin CJ: Effects of valsalva manoeuvre on maternal and fetal wellbeing. *Br J Midwifery* 2009; 17: 279–285
- 29 Brancato RM, Church S, Stone PW: A meta-analysis of passive descent versus immediate pushing in nulliparous women with epidural analgesia in the second stage of labor. *J Obstet Gynecol Neonat Nurs* 2008; 37: 4–12
- 30 de Tayrac R, Letouzey V: Methods of pushing during vaginal delivery and pelvic floor and perineal outcomes: a review. *Curr Opin Obstet Gynecol* 2016; 28: 470–476
- 31 Cooke A: When will we change practice and stop directing pushing in labour? *Br J Midwifery* 2010; 18: 76–81
- 32 WHO Reproductive Health Library: WHO recommendation on definition and duration of the second stage of labour. World Health Organization 2018
- 33 World Health Organization: WHO recommendations on intrapartum care for a positive childbirth experience. World Health Organization 2018
- 34 WHO Reproductive Health Library: WHO recommendation on fundal pressure to facilitate childbirth. World Health Organization 2018
- 35 Lehmann V: Warum denn immer ziehen und warum nicht auch mitunter drücken? *Z Gebursh Neonatol* 2017; 221: 247–248
- 36 Schoch F, Sebök N: Schneller mit Kristeller: eine historische Recherchearbeit zum Kristeller-Handgriff im Vergleich mit aktueller Literatur. Bachelor-Thesis an der Zürcher Hochschule für Angewandte Wissenschaften, 2016. DOI: 10.21256/zhaw-1220
- 37 Knabl J, Knüppel M, Karl K et al.: Wie lang ist zu lang?
– Eine Analyse der protrahierten Austreibungsperiode im Verhältnis zum Entbindungsmodus. *Gebursh Frauenheilk* 2008; 68 (S 01): PO_Geb_02_24
- 38 Cheng YW, Shaffer BL, Nicholson JM, Caughey AB: Second stage of labor and epidural use: a larger effect than previously suggested. *Obstet Gynecol* 2014; 123: 527–535
- 39 WHO Reproductive Health Library: WHO recommendation on techniques for preventing perineal trauma in second stage of labour. World Health Organization 2018
- 40 Sackett DL: Evidence-based medicine. *Semin Perinatol* 1997; 21: 3–5
- 41 Straus SE, Glasziou P, Richardson WS, Haynes RB: Evidence-Based Medicine E-Book: How to Practice and Teach EBM. Elsevier Health Sciences 2018