

Schmerzmanagement in Zeiten der Gehirnentwicklung

Hebammenforum 2013; 14: 1023-1029

- 1 Anand KJ, Hall RW: Love, pain, and intensive care. *Pediatrics* 2008; 121: 825-827
- 2 Damm G: Frühe Vergleiche: Wie entwickeln sich fünfjährige ehemalige Frühgeborene im Vergleich zu gleichaltrigen Reifgeborenen? *Niedersächsisches Ärzteblatt* 09/2012; S. 28-29
- 3 Grunau RE, Holsti L, Peters JW: Long-term consequences of pain in human neonates. *Semin Fetal Neonat Med* 2006; 11: 268-275
- 4 Petrack EM et al.: Pain Management in the Emergency Department: Patterns of Analgesic Utilization *PEDIATRICS* Vol. 99 No. 5 May 1997, pp. 711-714
- 5 Bauchner H et al.: Use of analgesic agents for invasive medical procedures in pediatric and neonatal intensive care units *The Journal of Pediatrics*, Volume 121, Issue 4, October 1992, Pages 647-649
- 6 Wolfe J: Suffering in children at the end of life: recognizing an ethical duty to palliate. *J Clin Ethics* 2000; 11(2): 157e163
- 7 Bremerich DH et al.: Postoperative Schmerztherapie im Kindesalter. Ergebnisse einer repräsentativen Umfrage in Deutschland. *Anästhesist* 2001; 50: 102-112
- 8 Sittl R et al.: Postoperative Schmerztherapie bei Kindern und Jugendlichen. *Schmerz* 2000; 14: 333-339
- 9 Gharavi B et al.: Pain management and the effect of guidelines in neonatal units in Austria, Germany and Switzerland. *Pediatr Int*. 2007; 49: 652-658
- 10 Anand KJ: Clinical importance of pain and stress in preterm neonates. *Biol Neonate* 1998; 73: 1-9
- 11 Grunau R: Early pain in preterm infants. A model of long-term effects. *Clin Perinatol* 2002; 29: 373-394
- 12 Gressens P et al.: The impact of neonatal intensive care practices on the developing brain. *J Pediatr* 2002; 140: 646-653
- 13 Berardi N, Pizzorusso T, Maffei L: Critical periods during sensory development. *Curr Opin Neurobiol* 2000; 10: 138-145
- 14 Hermann C et al.: Long-term alteration of pain sensitivity in school-aged children with early pain experiences. *Pain* 2006; 125: 278-285
- 15 Twycross A: Managing pain during the first year of life. *Infant* 2006; 2: 203-207
- 16 Taddio A, Katz J: The effects of early pain experience in neonates on pain responses in infancy and childhood. *Paediatr Drugs* 2005; 7: 245-257
- 17 Mayes Let al.: Attention regulation in infants born at risk. *Prematurity and Prenatal Cocaine Exposure* In: Burack, Enns (Hrsg.): *Attention, Development and Psychopathology*. Guilford Press, NYC 1997 ISBN 1-57230-198-8
- 18 D'Apolito KC: State of the science: procedural pain management in the neonate. *J Perinat Neonat Nurs* 2006; 20: 56-61
- 19 Leef KH: Evidence-based review of oral sucrose administration to decrease the pain response in newborn infants. *Neonat Netw* 2006; 25: 275-284
- 20 Stevens B et al.: Consistent management of repeated procedural pain with sucrose in preterm neonates: is it effective and safe for repeated use over time? *Clin J Pain* 2005; 21: 543-548
- 21 Holsti L et al.: Behavioral responses to pain are heightened after clustered care in preterm infants born between 30 and 32 weeks gestational age. *Clin J Pain* 2006; 22: 757-764
- 22 Holsti L et al.: Relationships between adrenocorticotrophic hormone and cortisol are altered during clustered nursing care in preterm infants born at extremely low gestational age. *Early Hum Dev* 2007; 83: 341-348
- 23 Holsti L et al.: Prior pain induces heightened motor responses during clustered care in preterm infants in the NICU. *Early Hum Dev* 2005; 81: 293-302
- 24 Deutsches Netzwerk für Qualitätsentwicklung in der Pflege (DNQP) 2003, www.wiso.hs-osnabrueck.de/38089.html (1. Aktualisierung Dezember 2011)
- 25 www.awmf.org/leitlinien/detail/ll/001-012.html, Stand: 1.12.09, gültig bis 1.12.14
- 26 Shah VS, Ohlsson A: Venepuncture versus heel lance for blood sampling in term neonates. *Cochrane Database Syst Rev*. 2011 Oct 5; (10): CD001452. doi: 10.1002/14651858.CD001452.pub4.
- 27 Cignacco E et al.: The efficacy of non-pharmacological interventions in the management of procedural pain in preterm and term neonates. A systematic literature review. *Eur J Pain* 2007; 11: 139-152
- 28 Stoffel L et al.: The efficacy of non-pharmacological interventions in the management of pain in preterm and term neonates. A systematic literature review. *Pflege* 2005; 18: 147-158
- 29 Akcan E, Yiğit R, Atıcı R: The effect of kangaroo care on pain in premature infants during invasive procedures. *Turk J Pediatr* 2009; 51: 14-18
- 30 Acharya AB et al.: Oral sucrose analgesia for preterm infant venipuncture. *Arch Dis Child Fetal Neonatal Ed* 2004; 89: F17-F18
- 31 Chermont AG et al.: Skin-to-skin contact and/or oral 25% dextrose for procedural pain relief for term newborn infants. *Pediatrics* 2009; 124: e1101-1107
- 32 Gray L, Wyatt L, Blass EM: Skin to skin contact is analgesic in healthy newborns. *Pediatrics* 2000; 105: 14-16
- 33 Axelin A, Salanterä S, Lehtonen L: »Facilitated tucking by parents« in pain management of preterm infants: a randomised crossover trial. *Early Hum Dev* 2006; 82: 241-247
- 34 Hill S et al.: Effects of facilitated tucking during routine care of infants born preterm. *Pediatr Phys Ther* 2005; 17: 158-163
- 35 Huang CM et al.: Comparison of pain responses of premature infants to the heelstick between containment and swaddling. *J Nurs Res* 2004; 12: 31-40
- 36 Arora J, Kumar, Ramji S: Effect of oil massage on growth and neurobehavior in very low birth weight preterm neonates. *Indian Pediatr* 2005; 42: 1092-1100
- 37 Ludington-Hoe SM, Hosseini R, Torowicz DL: Skin-to-skin contact (kangaroo care) analgesia for preterm infant heel stick. *AACN Clin Issues* 2005; 16: 373-387
- 38 Cong X et al.: Effects of Skin-to-Skin Contact on Autonomic Pain Responses in Preterm Infants *The Journal of Pain*, Vol 13, No 7 (July), 2012: pp 636-64
- 39 Diego MA, Field T, Hernandez-Reif M: Vagal activity, gastric motility, and weight gain in massaged preterm neonates. *J Pediatr* 2005; 147: 50-55
- 40 Beachy JM: Premature infant massage in the NICU. *Neonatal Netw* 2003; 22: 39-45
- 41 Weller A, Feldman R: Emotion regulation and touch in infants: the role of cholecystokin and opioids. *Peptides* 2003; 24: 779-788
- 42 Arnon S et al.: Live music is beneficial to preterm infants in the neonatal intensive care unit environment. *Birth* 2006; 33: 131-136
- 43 Chou LL et al.: Effects of music therapy on oxygen saturation in premature infants receiving endotracheal suctioning. *J Nurs Res* 2003; 11: 209-216
- 44 Standley JM: A meta-analysis of the efficacy of music therapy for premature infants. *J Pediatr Nurs* 2002; 17: 107-113
- 45 Carbajal R et al.: Analgesic effect of breast feeding in term neonates: randomised controlled trial. *BMJ* 2003; 326: 13
- 46 Gray L et al.: Breastfeeding is analgesic in healthy newborns. *Pediatrics* 2002; 109: 590-593
- 47 Shah PS, Aliwalas LI, Shah V: Breastfeeding or breast milk for procedural pain in neonates. *Cochrane Database Syst Rev* 2006; (3): CD004950
- 48 Mörelius E et al.: Is a nappy change stressful to neonates? *Early Hum Dev* 2006; 82: 669-676