

»Nehmt uns nicht auch noch den Dammschutz weg«

Hebammenforum 2014; 15: 226-228

- 1 Aasheim V et al.: Perineal techniques during the second stage of labour for reducing perineal trauma. *Cochrane database Syst Rev* 2011, CD006672
Zu finden unter www.ncbi.nlm.nih.gov/pubmed/22161407
- 2 Kindberg S, Seehafer P: *Geburtsverletzungen – vermeiden, erkennen, versorgen*. 1. Auflage Hannover: Elwin Staude, 2013
- 3 Dannecker C et al.: Episiotomy and perineal tears presumed to be imminent: randomized controlled trial. *Acta Obstet Gynecol Scand* 2004; 83: 364–368. Zu finden unter www.blackwell-synergy.com/links/doi/10.1111/j.0001-6349.2004.00366.x
- 4 Hartmann K et al.: Outcomes of routine episiotomy: a systematic review. *JAMA* 2005; 293: 2141–2148. Zu finden unter <http://jama.jamanetwork.com/article.aspx?articleid=200799>
- 5 Klein MC et al.: Does episiotomy prevent perineal trauma and pelvic floor relaxation? *Online J Curr Clin Trials* 1992; Doc No 10: [6019 words; 65 paragraphs]. Zu finden unter <http://europepmc.org/abstract/MED/1343606>
- 6 Klein MC et al.: Relationship of episiotomy to perineal trauma and morbidity, sexual dysfunction, and pelvic floor relaxation. *Am J Obstet Gynecol* 1994; 171: 591–598. Zu finden unter www.sciencedirect.com/science/article/pii/S0002937894900701
- 7 Lede RL, Belizán JM, Carroli G: Is routine use of episiotomy justified? *Am J Obstet Gynecol* 1996; 174: 1399–1402. Zu finden unter www.sciencedirect.com/science/article/pii/S0002937896705793
- 8 Poen AC et al.: Third degree obstetric perineal tears: risk factors and the preventive role of mediolateral episiotomy. *BJOG An Int J Obstet Gynaecol* 1997; 104: 563–566. Zu finden unter <http://doi.wiley.com/10.1111/j.1471-0528.1997.tb11533.x>
- 9 EURO-PERISTAT Project with SCPE and EUROCAT: *European Perinatal Health Report. The health and care of pregnant women and babies in Europe in 2010*. 2013. Zu finden unter <http://europeristat.com/reports/european-perinatal-health-report-2010.html>
- 10 AQUA: 16/1 – Geburtshilfe. Qualitätsindikatoren. Bundesauswertung zum Erfassungsjahr 2012. Göttingen, 2013. Zu finden unter www.sqg.de
- 11 Ghionzoli M et al.: Human amniotic fluid stem cell differentiation along smooth muscle lineage. *FASEB J* 2013; 27: 4853–4865. Zu finden unter www.fasebj.org/content/27/12/4853.short
- 12 Cluett ER, Burns E: Immersion in water in labour and birth. *Cochrane database Syst Rev* 2009; CD000111. Zu finden unter www.ncbi.nlm.nih.gov/pubmed/19370552
- 13 Reed R, Peetze J, Barnes M: Birth: an evidence-based approach. *J Paramed Pract* 2010; 2: 6–11. Zu finden unter www.paramedicpractice.com/cgi-bin/go.pl/library/abstract.html?uid=46148
- 14 Laine K et al.: Incidence of obstetric anal sphincter injuries after training to protect the perineum: cohort study. *BMJ Open* 2012; 2. Zu finden unter www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3488722&tool=pmcentrez&rendertype=abstract
- 15 Hals E et al.: A Multicenter Interventional Program to Reduce the Incidence of Anal Sphincter Tears. *Obstet Gynecol* 2010; 116. Zu finden unter http://journals.lww.com/greenjournal/Fulltext/2010/10000/A_Multicenter_Interventional_Program_to_Reduce_the.16.aspx
- 16 Fretheim A et al.: The impact of an intervention programme employing a hands-on technique to reduce the incidence of anal sphincter tears: interrupted time-series reanalysis. *BMJ Open* 2013; 3: e003355. Zu finden unter <http://bmjopen.bmj.com/content/3/10/e003355.short>
- 17 Schwarz C: Halten oder nicht halten? *Hebammenforum* 2009; 10: 269
- 18 Schwarz C: Dammschutz zwischen Tun und Lassen. *Dtsch Hebammenzeitschrift* 2011; 8: 6–9
- 19 Schwarz C: Jeder Dammschutz ist eine Intervention. *Dtsch Hebammenzeitschrift* 2008; 11: 17–20