

AUFMERKSAM UND GEDULDIG – TEIL 1: ÄNGSTE

Hebammenforum 3/2021; 22: 44–46

- 1 Dührssen A: Geburtshülfliches Vademecum für Studierende und Ärzte. Band 1. Karger 1908
- 2 Chalubinski KM: Normale Geburt. In: Schneider H, Husslein P, Schneider KTM: Die Geburtshilfe. Springer 2016: 663–686
- 3 Intrapartum care for healthy women and babies. National Institute for Health and Care Excellence: Clinical Guidelines. 2014
- 4 Care in second stage of labour. National Institute for Health and Care Excellence: Pathways. 2014
- 5 Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen (IQWiG): V16-01A – Definitionen der Geburtsphasen – Rapid Report – Version 1.0. 2017
- 6 Janni W, Schiessl B, Peschers U et al.: The prognostic impact of a prolonged second stage of labor on maternal and fetal outcome. *Acta Obstet Gynecol Scand* 2002; 81: 214–221
- 7 El-Hamamy E, Arulkumaran S: Poor progress of labour. *Curr Obstet Gynaecol* 2005; 15: 1–8
- 8 Lee N, Firmin M, Gao Y, Kildea S: Perineal injury associated with hands on/hands poised and directed/undirected pushing: A retrospective cross-sectional study of non-operative vaginal births, 2011–2016. *Int J Nurs Stud* 2018; 83: 11–17
- 9 Garmi G, Peretz H, Braverman M et al.: Risk factors for obstetric anal sphincter injury: To prolong or to vacuum? *Midwifery* 2016; 34: 178–182
- 10 Soong B, Barnes M: Maternal position at midwife-attended birth and perineal trauma: is there an association? *Birth* 2005; 32: 164–169
- 11 Eason E, Labrecque M, Wells G, Feldman P: Preventing perineal trauma during childbirth: a systematic review. *Obstet Gynecol* 2000; 95: 464–471
- 12 East CE, Lau R, Biro MA: Midwives' and doctors' perceptions of their preparation for and practice in managing the perineum in the second stage of labour: a cross-sectional survey. *Midwifery* 2015; 31: 122–131
- 13 Smith LA, Price N, Simonite V, Burns EE: Incidence of and risk factors for perineal trauma: a prospective observational study. *BMC Pregnant Childbirth* 2013; 13: 1–9
- 14 Cortes E, Basra R, Kelleher CJ: Waterbirth and pelvic floor injury: a retrospective study and postal survey using ICIQ modular long form questionnaires. *Eur J Obstet Gynecol Reprod Biol* 2011; 155: 27–30
- 15 Memon HU, Handa VL: Vaginal childbirth and pelvic floor disorders. *Womens Health* 2013; 9: 265–277
- 16 Danforth KN, Townsend MK, Lifford K et al.: Risk factors for urinary incontinence among middle-aged women. *Am J Obstet Gynecol* 2006; 194: 339–345
- 17 Minassian VA, Stewart WF, Wood GC: Urinary incontinence in women: variation in prevalence estimates and risk factors. *Obstet Gynecol* 2008; 111: 324–331
- 18 Luber KM: The definition, prevalence, and risk factors for stress urinary incontinence. *Rev Urol* 2004; 6 (Suppl 3): S3–9
- 19 Kearney R, Miller JM, Ashton-Miller JA, DeLancey JO: Obstetric factors associated with levator ani muscle injury after vaginal birth. *Obstet Gynecol* 2006; 107: 144–149
- 20 Dietz H, Steensma AB: The role of childbirth in the aetiology of rectocele. *BJOG* 2006; 113: 264–267
- 21 Kopas ML: A review of evidence-based practices for management of the second stage of labor. *J Midwifery Womens Health* 2014; : 264–276
- 22 www.who.int/reproductivehealth/topics/maternal_perinatal/fistula/en (Zugriff 21.10.20)
- 23 De Bernis L: Obstetric fistula: guiding principles for clinical management and programme development, a new WHO guideline. *Int J Gynecol Obstet* 2007; 99: S117–S121
- 24 Tunçalp Ö, Tripathi V, Landry E, Stanton CK, Ahmed S: Measuring the incidence and prevalence of obstetric fistula: approaches, needs and recommendations. *Bull World Health Organ* 2015; 93, 60–62
- 25 Dahlen HG, Homer CSE, Leap N, Tracy SK: From social to surgical: historical perspectives on perineal care during labour and birth. *Women Birth* 2011; 24: 105–111