

KLEINE UND MITTLERE GEBURTSVERLETZUNGEN VERMEIDEN

Hebammenforum 4/2022; 23: 8–14

- 1 WHO: International Classification of Diseases 11th Revision, 2022, <https://icd.who.int/en>
- 2 IQTIG-Institut (Hrsg.): Bundesauswertung der klinischen Geburtshilfe in Deutschland 2021. Erfassungsjahr 2020
- 3 Beckmann MM, Stock OM: Antenatal perineal massage for reducing perineal trauma. Cochrane Database Syst Rev 2013; Issue 1. Art. No.: CD005123
- 4 World Health Organization Division of Family Health Maternal Health and Safe Motherhood: Care in normal birth: a practical guide. Report of a technical working group. World Health Organization, Geneva 1996
- 5 Hillebrenner J et al.: Erste klinische Erfahrungen bei Erstgebärenden mit einem neuartigen Geburtstrainer Epi-No®. Z Geburtshilfe Neonatol 2001; 205: 12–19
- 6 Kamisan Atan I et al.: Does the Epi-No® birth trainer prevent vaginal birth-related pelvic floor trauma? A multicentre prospective randomised controlled trial. BJOG 2016; 123 (6): 995–1003
- 7 Rodríguez-Blanque R et al.: Randomized Clinical Trial of an Aquatic Physical Exercise Program During Pregnancy. J Obstet Gynecol Neonatal Nurs 2019 May; 48 (3): 321–331
- 8 Carrascosa MDC et al.: Aquanatal Trial. Effect of aerobic water exercise during pregnancy on epidural use and pain: A multi-centre, randomised, controlled trial. Midwifery 2021 Dec; 103: 103105
- 9 Leon-Larios F et al.: Influence of a pelvic floor training programme to prevent perineal trauma: A quasi-randomised controlled trial. Midwifery 2017 Jul; 50: 72–77
- 10 Sobhgal SS et al.: The effect of antenatal pelvic floor muscle exercises on labour and birth outcomes: a systematic review and meta-analysis. Int Urogynecol J 2020; 31: 2189–2203
- 11 Zabados S: Auswirkungen der geburtsvorbereitenden präpartalen Akupunktur auf das Geburtsergebnis bei Erst- und Mehrgebärenden. Dissertation zur Erlangung des Grades eines Doktors der Medizin. Hamburg 2009, Universität Hamburg
- 12 Caudwell-Hall J et al.: Predictors of maternal levator ani injury. Acta Obstet Gynecol Scand 2017 Apr; 96 (4): 426–431
- 13 Aiken CE et al.: Influence of the duration of the second stage of labor on the likelihood of obstetric anal sphincter injury. Birth 2015; 42 (1): 86–93
- 14 Runge M: Lehrbuch der Geburtshilfe. Springer Verlag, Berlin 1903: 142
- 15 Samuelsson E et al.: A prospective observational study on tears during vaginal delivery: occurrences and risk factors. Acta Obstet Gynecol Scand 2002; 81: 44–49
- 16 Fitzgerald MP et al.: Pelvic Floor Disorders Network: Risk factors for anal sphincter tear during vaginal delivery. Obstet Gynecol 2007, 109 (1): 29–34
- 17 Meister MR et al.: Predicting obstetric anal sphincter injuries in a modern obstetric population. Am J Obstet Gynecol 2016, 215 (3): 310
- 18 Hastings-Tolsma M et al.: Getting through birth in one piece: protecting the perineum. MCN Am J Matern Child Nurs 2007, 32 (3): 158–164
- 19 Wright A et al.: Safe Motherhood and Newborn Health Committee. FIGO good clinical practice paper: management of the second stage of labor. Int J Gynaecol Obstet 2021 Feb; 152 (2): 172–181
- 20 Sveinsdóttir E et al.: Effects of an intervention program for reducing severe perineal trauma during the second stage of labor. Birth 2019 Jun; 46 (2): 371–378
- 21 Cluett ER et al.: Immersion in water during labour and birth. Cochrane Database Syst Rev 2018 May 16; 5 (5): CD000111
- 22 Sidebottom AC et al.: Maternal and Neonatal Outcomes in Hospital-Based Deliveries With Water Immersion. Obstet Gynecol 2020 Oct; 136 (4): 707–715
- 23 Neiman E et al.: Outcomes of Waterbirth in a US Hospital-Based Midwifery Practice: A Retrospective Cohort Study of Water Immersion During Labor and Birth. J Midwifery Womens Health 2020 Mar; 65 (2): 216–223
- 24 Aasheim V et al.: Perineal techniques during the second stage of labour for reducing perineal trauma. Cochrane Database of Syst Rev 2017, Issue 6. Art. No.: CD006672
- 25 Aquino CI et al.: Perineal massage during labor: a systematic review and meta-analysis of randomized controlled trials. J Matern Fetal Neonatal Med 2020 Mar; 33 (6): 1051–1063
- 26 Magoga G et al.: Warm perineal compresses during the second stage of labor for reducing perineal trauma: A meta-analysis. Eur J Obstet Gynecol Reprod Biol 2019 Sep; 240: 93–98
- 27 Stiefel A, Geist C, Harder U (Hrsg.): Hebammenkunde. Lehrbuch für Schwangerschaft, Geburt, Wochenbett und Beruf. 5. überarb. und erw. Aufl., Hippokrates Verlag, Stuttgart 2013
- 28 Hilde G et al.: How well can pelvic floor muscles with major defects contract? A cross-sectional comparative study 6 weeks after delivery using transperineal 3D/4D ultrasound and manometer. BJOG 2013, 120 (11): 1423–9
- 29 Aasheim V et al.: Perineal techniques during the second stage of labour for reducing perineal trauma. Cochrane Database Syst Rev 2017, Issue 6. Art. No.: CD006672
- 30 Laine K et al.: Incidence of obstetric anal sphincter injuries after training to protect the perineum: cohort study. BMJ open. 2 (5): e001649
- 31 Bulchandani S et al.: Manual perineal support at the time of childbirth: a systematic review and meta-analysis. BJOG 2015; 122 (9): 1157–65
- 32 Rasmussen OB et al.: Reducing the incidence of Obstetric Sphincter Injuries using a hands-on technique: an interventional quality improvement project. BMJ Qual Improv Rep 2016; 5: 1
- 33 Lemos A et al.: Pushing/bearing down methods for the second stage of labour. Cochrane Database Syst Rev. 2017 Mar 26; 3 (3): CD009124
- 34 Araújo AE et al.: Efficacy of spontaneous pushing with pursed lips breathing compared with directed pushing in maternal and neonatal outcomes. J Obstet Gynaecol 2021 Sep 28: 1–7
- 35 Aabakke AJ et al.: The effect of maneuvers for shoulder delivery on perineal trauma: a randomized controlled trial. Acta Obstet Gynecol Scand 2016 Sep; 95 (9): 1070–7
- 36 Xiao L et al.: A core outcome set for clinical trials of first- and second-degree perineal tears prevention and treatment: a study protocol for a systematic review and a Delphi survey. Trials 2021; 22: 843